



Contribution ID: 93

Type: Oral Presentation

Linear polarization measurement on gamma rays from non-oriented nuclear states

Tuesday 8 July 2025 09:20 (20 minutes)

Author : Beatrice Similindi

Co-author : Dr E.A Lawrie

Dr S.N.T Majola

Gamma ray spectroscopy is a powerful tool in nuclear structure, but the interpretation of the intrinsic properties of the nucleus becomes complex if the parity of the emitted radiation is not known. This study aimed at developing a technique to measure linear polarization of γ rays emitted

from non-oriented nuclear states for the clover detectors of the iThemba LABS AFRODITE and GAMKA arrays. Orientation was created by gating on a γ ray detected in one clover detector while observing another γ ray detected in coincidence in another detector. Having being well known for

their high efficiency in detecting γ rays, the clover detectors used in this study comprise 4 Ge crystals which were used as Compton polarimeter, which allows us to measure linear polarization. A focus was made on studying the Compton scattering of γ rays across the 4 Ge crystals in which the Klein-Nishina equation was utilised for the two experimental set-ups to measure the polarization anisotropy. Theoretical curves for all observed rays emitted from the beta-decay of ^{152}Eu were derived and compared with the measured polarization anisotropy, A_p to deduce the polarization sensitivity, $Q(\gamma)$. Furthermore, the degree of linear polarization was determined experimentally for the γ rays observed in ^{196}Hg following the beta-decay of ^{196}Tl using the deduced polarization sensitivity for the AFRODITE array. The technique has also been applied

for the upgraded GAMKA array. The polarization sensitivity for the GAMKA array was determined using experimental data with Eu source by measuring the linear polarization anisotropy and comparing it with the theoretical curves for linear polarization for the well-known pure transitions.

The derived linear polarization sensitivity Q was then used to determine mixing ratios of observed mixed transitions. In addition, the angular correlation coefficients for both pure and mixed transition have been measured and compared with the theoretical coefficients. Lastly, final values for the mixing ratios for $M1+E2$ and $E0+M1+E2$ transitions have been determined by combining angular correlation and linear polarization results for all observed γ ray cascades detected with the GAMKA

array and with Eu source. The deduced polarization sensitivity can be used in future experiments that intend to measure γ rays whose parities as well as mixing ratios are not known.

[1] K. Krane, Introductory Nuclear Physics, chapters 7.1 (Electro Magnetic Radiation), 7.9 (Polarimeters), 10.1, 10.2, (optional 10.3), 10.4, 10.5.

[2] G. Duchene et al., Nucl. Instrum. Meth. A, 432, 90 (1999).

[3] Fagg, L. W. and Hanna, S. S. Polarization measurements on nuclear γ - rays. Reviews of Modern Physics, 31(3):711, 1959.

[4] P. Jones et al., Nucl. Instrum. Meth. A, 362, 556 (1995).

[5] A. Wolf et al., Phys. Rev. C 66, 024323 (2002).

Apply for student award at which level:

MSc

Consent on use of personal information: Abstract Submission

Yes, I ACCEPT

Primary author: Ms SIMILINDI, Beatrice (iThemba LABS)

Co-authors: LAWRIE, Elena Atanassova (iThemba LABS); LAWRIE, Jacobus (University of the Western Cape); MTHEMBU, Sinegugu (University of the Western Cape & iThemba LABS); MAJOLA, Siyabonga (University of Johannesburg)

Presenter: Ms SIMILINDI, Beatrice (iThemba LABS)

Session Classification: Nuclear, Particle and Radiation Physics-1

Track Classification: Track B - Nuclear, Particle and Radiation Physics