



Contribution ID: 502

Type: Oral Presentation

Radiation contamination in gold mine tailings soil samples using HPGe spectrometry

Tuesday 8 July 2025 14:40 (20 minutes)

An assessment of the radiation concentration in abandoned mines located near settlements in the west of Johannesburg was carried out, and the corresponding radiological indices were determined. In this study, a gamma spectrometer was used to measure the activity concentrations of radionuclides in the soil samples. The activity concentrations for ^{226}Ra , ^{232}Th , and ^{40}K were 338.44 ± 3.48 , 10.06 ± 0.68 , and 126.15 ± 10.90 Bq/kg, respectively. The results revealed that the average activity concentrations at some locations exceeded the world average for some of these nuclides.

Apply for student award at which level:

Consent on use of personal information: Abstract Submission

Yes, I ACCEPT

Primary authors: Dr KHESWA, B.V. (University of Johannesburg); Prof. MASITENG, L.P. (University of Johannesburg); MVELASE, M.J.; Dr MALEKA, P.P. (iThemba Labs); Prof. NTSHANGASE, S.S. (University of Zululand); Dr MAJOLA, Siyabonga (University of Johannesburg)

Presenter: MVELASE, M.J.

Session Classification: Nuclear, Particle and Radiation Physics-1

Track Classification: Track B - Nuclear, Particle and Radiation Physics